



BIOSTAR TOP

SPECIFIC ACTION PRODUCT
Inoculation of mycorrhizal fungi

ALLOWED IN ORGANIC FARMING



BIOAGRONOMIC CHARACTERISTICS

BIOSTAR TOP contains microorganisms and noble milk proteins which, by rebalancing the microbial flora present in the environment and on the plant, allow optimal development of vegetative growth, positively predisposing to overcome all abiotic and biotic stresses.

FINAL RESULT

- Improvement of the state of well-being of the plant.
- Reduction of stress from pathogenic attacks.
- Its nutrition helps to overcome the stresses of White Disease
- Fruit and vegetables without residue.

DOSES AND METHODS OF USE

Type of application: FOLIAR

It can be used on all crops:

VINE: 2.5 kg/ha

TREES: (Pome fruit, stone fruit, Actinidia, Citrus and Olive trees) 2.5-4 kg/ha

HORTICULTURAL: 2.5-4 kg/ha

STRAWBERRY AND SMALL FRUITS: 2.5-3 kg/ha

NURSERIES: 2-4 kg/ha

GREENHOUSE: 250-400gr/1000sqm

APPLICATIONS • *It can be used from the beginning of the cycle to mature with a frequency of 8 days, if necessary, in hot and humid periods reduce to 5 days.*

It has no shortage times and leaves no residue.

WARNINGS • *The product is miscible with most pesticides other than products containing copper or antibacterials. However, you should take a test before applying. The manufacturer is not liable for damage caused by improper use of the product.*

COMPOSITION

Type of organic soil improver:
simple non-composted
vegetable soil improver

- Mycorrhizal content 0.1%
- Rhizosphere bacteria content 10^7 CFU/gr
- Non-genetically modified lactic acid bacteria 10^7 CFU/gr

FORMULATION LIQUID

PACKAGING

1 kg bottle
5 kg tank
10 kg tank

SIAN Reg. No.
IT 11423/15

CERT. CCPB
N° 05/2018 MT



SPECIALAGRI S.R.L.

Via Veneto 8 • Cologna Veneta (VR) • Italy

Tel. +39 0442 412833

info@specialagri.it

www.specialagri.it